




January 1 – November 4, 2017

Epidemiology Bureau
Public Health Surveillance Division

TRENDS IN THE PHILIPPINES

This report summarizes the Influenza-like Illness (ILI) surveillance activities nationwide. A total of **107,689** Influenza-like Illness cases were reported nationwide from January 1 to November 4, 2017. This is **11.49%** lower compared to the same time period last year (**121,674**). For Influenza cases, a total of **211** confirmed cases were reported nationwide from January 1 to November 4, 2017. This is **17.58%** lower compared to the same time period last year (**256**).

Fig. 1 Confirmed Influenza Cases by Morbidity Week 2017* vs 2016, Philippines, as of January 1 - November 4, 2017 (N=211)



GEOGRAPHIC DISTRIBUTION

The number of confirmed influenza cases vary by region. Table 1 shows the distribution of reported confirmed influenza cases and deaths by region in comparison to the same time period last year. There is an increase in the number of reported cases in all regions. No reported deaths among the confirmed influenza cases.

Majority of the reported confirmed influenza cases came from NCR (78 cases, 36.97%), CAR (43 cases, 20.38%) and Region V (33 cases, 15.64%) as shown in Figure 2.


Table 1. Confirmed Influenza Cases by Region 2017* vs 2016, Philippines, as of January 1 - November 4, 2017 (N=211)

Fig. 2 Confirmed Influenza Cases by Region, Philippines, as of January 1 - November 4, 2017 (N=211)

| Region | Cases | | | Deaths | | | |
|-------------|-------|------|------------|--------|---------|------|---------|
| | 2017 | 2016 | % Change | 2017 | CFR (%) | 2016 | CFR (%) |
| I | 0 | 0 | ⇒ 0.00 | 0 | 0.00 | 0 | 0.00 |
| II | 0 | 0 | ⇒ 0.00 | 0 | 0.00 | 0 | 0.00 |
| III | 1 | 0 | ↑ 100.00 | 0 | 0.00 | 0 | 0.00 |
| IV-A | 0 | 0 | ⇒ 0.00 | 0 | 0.00 | 0 | 0.00 |
| MIMAROPA | 1 | 0 | ↑ 100.00 | 0 | 0.00 | 0 | 0.00 |
| V | 33 | 16 | ↑ 106.25 | 0 | 0.00 | 0 | 0.00 |
| VI | 0 | 0 | ⇒ 0.00 | 0 | 0.00 | 0 | 0.00 |
| VII | 13 | 0 | ↑ 1,300.00 | 0 | 0.00 | 0 | 0.00 |
| VIII | 0 | 0 | ⇒ 0.00 | 0 | 0.00 | 0 | 0.00 |
| IX | 31 | 83 | ↓ -62.65 | 0 | 0.00 | 0 | 0.00 |
| X | 0 | 0 | ↓ -200.00 | 0 | 0.00 | 0 | 0.00 |
| XI | 0 | 0 | ⇒ 0.00 | 0 | 0.00 | 0 | 0.00 |
| XII | 11 | 17 | ↓ -35.29 | 0 | 0.00 | 0 | 0.00 |
| ARMM | 0 | 0 | ⇒ 0.00 | 0 | 0.00 | 0 | 0.00 |
| CAR | 43 | 71 | ↓ -39.44 | 0 | 0.00 | 0 | 0.00 |
| CARAGA | 0 | 0 | ⇒ 0.00 | 0 | 0.00 | 0 | 0.00 |
| NCR | 78 | 69 | ↑ 13.04 | 0 | 0.00 | 0 | 0.00 |
| PHILIPPINES | 211 | 256 | ↓ -17.58 | 0 | 0.00 | 0 | 0.00 |

Legend - CONFIRMEDCASE

1 Dot = 1






January 1 – November 4, 2017

Epidemiology Bureau
Public Health Surveillance Division

PROFILE OF CASES

In Figure 3, there are 105 (49.76%) males and 106 (50.24%) females and majority of the confirmed influenza cases belonged to the 1 to 4 age group with 89 (42.18%) cases and median of 4 year old.


Fig. 3 Confirmed Influenza Cases by Age Group & Sex, Philippines, as of January 1 - November 4, 2017 (N=211)



INFLUENZA TYPE IN THE PHILIPPINES

Among the confirmed influenza cases, top Influenza types are as follows (Figure 4): *Influenza A/California/7/2009(H1N1)-like* (118 cases, 55.92%), *Influenza A/Perth/16/2009(H3N2)-like* (24 cases, 11.37%), *Influenza B/Yamagata/16/88-like* (15 cases, 7.11%) and *Influenza A* (11 cases, 5.21%).

Fig. 4 Organism of Confirmed Influenza Cases, Philippines, as of January 1 - November 4, 2017 (N=211)






January 1 – November 4, 2017

Epidemiology Bureau
Public Health Surveillance Division

Editorial Board


IRMA L. ASUNCION, MD, MHA, CESO IV
Director IV, Epidemiology Bureau


GENESIS MAY J. SAMONTE, MD, MSc, PHSAE
OIC-Division Chief, PHSD


MARIEL A. DEJESA, MD, MPM
Medical Officer IV, PHSD


JUNE CANTATA B. CORPUZ, RN
Nurse III, PIDS National Coordinator


JEZZA JONAH D. CRUCENA, RN
Nurse III, VPDS National Coordinator